

FULL PAPER

Cytotoxic activity of the purified extracts from duku (*Lansium domesticum* Corr.) Leaf against MCF-7 and HTB-183 cell lines, and the correlation with antioxidant activity

Muhammad Fauzan Lubis^{a,*} | Sumaiyah Sumaiyah^b | Embun Suci Nasution^c | Ririn Astyka^a | Masfria Masfria^d | Hafid Syahputra^d | Carlos Nofanolo Lase^a

^aDepartment of Pharmaceutical Biology, Universitas Sumatera Utara, Medan, 20222, Indonesia

^bDepartment of Pharmaceutical Technology, Universitas Sumatera Utara, Medan, 20222, Indonesia

^cDepartment of Pharmacology, Universitas Sumatera Utara, Medan, 20222, Indonesia

^dDepartment of Pharmaceutical Chemistry, Universitas Sumatera Utara, Medan, 20222, Indonesia

One of the plants that potential to develop as an anticancer agent is Duku leaf (*Lansium domesticum* Corr.). From previous studies, Duku leaf extract had cytotoxic activity against several cancerous cell lines. The aim of this study was to determine the cytotoxic activity of crude and purified extracts of Duku leaf against MCF-7 and HTB-183 cell lines. The crude extract of Duku leaf was obtained using maceration in ethanol absolute. The vacuum liquid chromatography with a gradient mobile phase was performed to obtain the purified extracts. The crude and purified extracts were observed to inhibit MCF-7 and HTB-183 cells using 3-(4,5-dimethyl thiazol 2-yl)-2,5-diphenyl tetrazolium bromide (MTT) assay. The antioxidant properties of samples were determined using radical DPPH scavenging activity. The data was collected and analyzed to provide the inhibitory concentration 50 (IC₅₀) and Pearson's correlation was analyzed to describe the correlation between cytotoxic and antioxidant properties. Five purified extracts were obtained and tested against MCF-7 and HTB-183. The purified extract D has stronger anticancer and antioxidant activities than other samples with IC₅₀ of 56.26 ± 3.11 µg/mL, 70.94 ± 2.92 µg/mL, and 53.65 ± 1.55 µg/mL, respectively (*p* < 0.05). We confirmed that cytotoxic activity has a strong correlation with antioxidant properties. Furthermore, the active compounds of purified extract D needed to be investigated and tested against MCF-7 and HTB-183 to explore the possible anticancer mechanisms.

*Corresponding Author:

Muhammad Fauzan Lubis

Email: fauzan.lubis@usu.ac.id

Tel.: +6281264353744

KEYWORDS

Lansium domesticum corr; vacuum liquid chromatography; cytotoxic; antioxidant.

Introduction

According to the World Health Organization (WHO), cancer is the primary cause of mortality on a global scale. In 2020, breast and lung cancer exhibited the highest incidence

rates, as reported by the WHO [1]. According to the American Society, it is projected that around 43,700 women will succumb to breast cancer in the United States in the year 2023 [2]. In addition, the estimated number of deaths from lung cancer is approximately

127,070, with 67,160 occurring in men and 59,910 in women [3]. China, as the largest and most densely populated upper-middle-income nation, has witnessed a notable increase in the incidence of breast cancer [4]. Breast cancer is the prevailing form of cancer, accounting for 16.6% of all cancer cases in Indonesia [5]. It stands as the second leading cause of cancer-related mortality, with a rate of 9.6%, following lung cancer, which holds a mortality rate of 13.2% [6].

The utilization of natural ingredients in cancer treatment has experienced a notable rise [7-9]. Herbal remedies have a strong historical and cultural significance in various societies and traditional practices [10-13]. There have been reports indicating that compounds derived from natural components had anticancer properties [14-17]. *Lansium domesticum* Corr., also referred to as Duku is a tropical fruit tree belonging to the Meliaceae family. It is extensively farmed primarily for local use and is indigenous to Southeast Asia, including Malaysia, Thailand, Indonesia, Vietnam, and the Philippines [18]. Multiple investigations have documented the potential medicinal qualities of Duku, including its anticancer [19], antidiarrheal [20], antipyretic [21], and anthelmintic [22] effects. The presence of alkaloids, flavonoids, tannins, triterpenoids/steroids, and saponins has been detected in the leaves of the Duku plant [18]. The primary objective of this investigation was to assess the anticancer properties of crude and purified extract from Duku leaves against MCF-7 and HTB-183 cell lines.

The MCF-7 cell line holds the distinction of being extensively investigated in human breast cancer research, making it the subject of several studies worldwide. The findings derived from this cell line have played a pivotal role in advancing our understanding of breast cancer and improving the prognosis and treatment options available to patients. Approximately 25,000 scientific articles have utilized this particular cell line. The MCF-7 cell line plays a significant role in elucidating the

efficacy of pharmaceutical agents for the treatment of breast cancer, particularly in the transition from non-metastatic to metastatic stages. Hence, MCF-7 cells continue to be employed for cancer medication development in contemporary research [23,24]. Meanwhile, similar to the exploration of anti-breast cancer agents, an *in vitro* study was conducted to find anti-lung cancer agents. Commonly, lung cancer is classified into two main groups, these are non-small cell lung cancer (NSCLC) and small cell lung cancer. The three major subtypes of NSCLC are squamous cell carcinoma, adenocarcinoma, and large cell carcinoma. These cancer types account for nearly 80% of all lung cancer cases. Therefore, to develop an anti-lung cancer drug approach, HTB-183 cells which are part of large cell carcinoma can be used because they are considered to reflect a frequently occurring lung cancer condition [25]. However, molecular characteristics of cancer cells regarding survival are a main factor that needs to be considered [26].

Several parts of Duku were used to assess the anticancer activity against various cancer cell lines. As reported by Manosroi *et al.* (2012), the parts like ripe fruits, stalks, and young fruits from Duku have activity to inhibit HT-29 and KB cell lines [19]. While, Khalili *et al.* (2017) were successful to compared the anticancer activity of several fruit extracts from Duku against HT-29 [21]. On the other hand, the focus on the anti-breast cancer activity of Duku parts, Fadhilah *et al.* (2020) explain the fruit peels of Duku exhibit cytotoxicity against the T47D cell line [27]. Recently, Fadhilah *et al.* (2020) reported that the fruit peels of Duku have anticancer activity against HepG2 [28]. A similar activity was reported by Lubis *et al.* (2022), but utilized the leaf part of Duku against HepG2 [29]. The utilization of Duku leaf as an anti-cancer not only against HepG2 cells. As reported by Lubis *et al.* (2023), the extract of Duku leaf has anti-pancreatic cancer against PANC-1 cells [30]. However, the study of Duku leaf extract and its

derived or fraction against MCF-7 and HTB-183 cells is not identified. This study was conducted to identify the anti-cancer activity of crude and purified extracts of Duku leaf against MCF-7 and HTB-183 cells. The process of purifying the extract involved the utilization of vacuum liquid chromatography. In addition, the mechanism by which purified extracts function as anticancer agents was elucidated through the assessment of their antioxidant activity utilizing the radical DPPH scavenging method.

Experimental

Materials

Dimethyl sulfoxide (DMSO), DMEM and RPMI medium, 0.25% trypsin EDTA, Fetal bovine serum, Fungizone® (Gibco, USA), 0.4% trypan blue, Penicillin-streptomycin, 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT reagent) (Sigma, USA), Ethanol, Ethyl acetate, and n-hexane (Merck, USA), silica gel 60 for column (0.063-0.200 mm) and 2,2-Diphenyl-1-picrylhydrazyl (DPPH) (Sigma, USA). The MCF-7 and HTB-183 cell lines were collected from the Parasitology Laboratory, Universitas Gadjah Mada, Yogyakarta, Indonesia.

Extract preparation and purification

Duku leaf (*Lansium domesticum* Corr.) was obtained in Medan, North Sumatera, Indonesia. The plant species were verified and identified by botanical practice in the Herbarium Medanense, Universitas Sumatera Utara, Medan, Indonesia (Voucher ID: 2084/HM-USU/12/2023). The leaves were washed, dried in a drying cabinet at 40-50 °C, and ground using the conventional blender. Extraction of dry powder was finished using maceration. The 180 g dry powder was put into a container, soaked with 1800 mL of solvent (ethanol absolute), and kept out for 24 h at room temperature. Afterward, the macerate was obtained after filtration using

Whatman No. 1. The macerate was evaporated using a rotary evaporator (Heidolph, Germany) to obtain a crude extract [31]. The crude extract was purified using vacuum liquid chromatography with silica gel 60 as an adsorbent and a combination of n-hexane: ethyl acetate in gradient (100:0 up to 10:90) and ethanol absolute for finishing the purification. The purified extracts were collected and categorized by Retention factor (Rf) values using thin-layer chromatography with sulfuric acid 50% as a visualization reagent. All purified extracts with the same kind of spotted profiles were grouped and kept in a temperature room before use [28,32].

Physicochemical analysis and phytochemical screening

Physicochemical analysis of dry powder and crude extract of Duku leaf include water content, total ash, acid insoluble ash content, water-soluble content, and ethanol soluble content [33]. Alkaloids were identified using color reagents such as Mayer, Bouchard, and Dragendroff. Shinoda and cyanidin test for flavonoids, tannin with FeCl₃ solution, and steroid/triterpenoids with Liebermann bouchard reagent [34].

Cytotoxic activity of crude and purified extract

The cytotoxicity activity was carried out using the MTT method. It is a calorimetric, non-radioactive, and quick test that converts 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT), which is produced when the mitochondrial enzymes in a healthy cell convert the yellow MTT substrate to the insoluble purple formazan [35]. MCF-7 and HTB-183 cells were planted into each well of 96-well plates. Afterward, varied concentrations of crude and purified extract including 500, 250, 125, 62,5, and 31,25 g.mL⁻¹, were applied to the planted cells, and incubated for 24 h at 37 °C. Excess media

was discarded and fresh media was added, then 10 μL of tetrazolium dye was added and incubated for 4 h. After incubation, a color change from yellow to purple was observed. MTT reaction stopped with stopper reagent (SDS 10% in 0.1 N HCl). In addition, the absorbances were detected with a microplate reader at 595 nm wavelength [36]. The % viability cell was measured based on control absorbance and sample absorbance.

Antioxidant activity of crude and purified extract

The measurement of free radical scavenging activity was conducted using the DPPH technique. A 0.2 mM solution of DPPH in methanol was produced. Subsequently, 100 μL of this solution was added to a solution containing extracts at a concentration of 100 $\mu\text{g}/\text{mL}$. After 60 minutes, the measurement of absorbance was conducted at a wavelength of 516 nm. The calculation of the percentage of inhibition was performed by comparing the absorbance values obtained from the control group with those obtained from the samples [37].

Statistical analysis

Statistical analyses were run on SPSS software version 22. The Kolmogorov-Smirnov test was used to check and confirm the normality of the

data distribution. Using the factorial ANOVA test, we evaluated the impact of the tested samples. Tukey's HSD post-hoc test was used to determine significance, and $p < 0.05$ was chosen as the threshold for statistical significance. Meanwhile, the correlation of cytotoxic and antioxidant activity was described using Pearson's correlation test.

Results and discussion

The determination of water content in the sample was set to maintain the quality of the sample. The less water content in dry powder and crude extract, the less possibility of contamination by mold growth. The water content should not be more than 10% for dry powder [38]. Determination of ash content was carried out to measure the total amount of external and internal minerals in a sample from the initial process to the end of manufacture. Determination of water and ethanol-soluble content was carried out to see the number of soluble compounds in water and ethanol solvents [39]. The results of the determination of dry powder and crude extract show that the results meet the requirements and are guaranteed quality based on *Materia Medika Indonesia* (MMI) [40] (Table 1).

TABLE 1 Physicochemical analysis of dry powder and crude extract of Duku leaf

Parameters	Dry powder	Crude extract
Water content (%)	6.62	10.55
Total ash content (%)	7.95	4.70
Acid-insoluble ash content (%)	4.45	2.52
Water soluble content (%)	18.40	NI
Ethanol soluble content (%)	12.60	NI

NI = Not Identified.

Based on the result of the phytochemical screening, Duku leaf has identified the presence of bioactive compounds and contains metabolites of alkaloid, flavonoid, saponin, steroid/triterpenoid, tannin, and glycoside. Dry powder and ethanol extract of

Duku leaf showed positive results containing alkaloid compounds in 3 tests characterized by the formation of a white residue after adding the Mayer reagent, a brown residue after the addition of Bouchardat reagent and no red-orange residue formed when the

Dragendorff reagent added [41]. From the results of the examination of flavonoids, the addition of concentrated hydrochloric acid to Mg powder and amyl alcohol formed a yellow-orange color layer on the amyl alcohol layer [42]. Compounds of the saponin group were declared to contain saponins due to the presence of stable foam after administration of hydrochloric acid [43]. Steroid/

triterpenoid examination showed positive steroid was indicated by the appearance of a green color. In the tannin test, a positive result is indicated by a change in the color of the filtrate to blackish-blue [44]. The result was supported by research that mentioned crude extract of Duku leaf obtained alkaloid, flavonoid, steroid/ triterpenoid, and tannin (Table 2) [18].

TABLE 2 Phytochemical screening of dry powder and crude extract of Duku leaf

Secondary metabolites	Dry powder	Crude extract
Alkaloids	Positive	Positive
Flavonoids	Positive	Positive
Saponin	Positive	Positive
Steroids/triterpenoids	Positive	Positive
Tannin	Positive	Positive
Glycoside	Positive	Positive

The crude extract was purified using vacuum liquid chromatography, yielding 10 fractions. Each fraction was grouped according to the profile of spotted after visualized using 50% sulfuric acid (Figure 1). The Rf values of spotted were identified and found in 5 purified extracts with different Rf values and color of the spot (Table 3). Two purified extracts, A and B come from one fraction each. Two spots were determined from purified extract A with Rf of 0.8125 (blue) and 0.875 (blue), and then three spots were identified from purified extract B with Rf 0.6875 (red), 0.8125 (red), and 0.875 (red), respectively. These results described the purified extracts A and B as almost similar based on the identified spots. The purified extract C was a purified extract from the merger of many fractions. Each fraction has 7 spots with Rf of 0.3125 (red), 0.4375 (red), 0.5625 (purple), 0.6875 (red), 0.8125 (red), and 0.875 (blue), respectively. In addition, the purified extract D was generated from a merger of two fractions. Fractions 8 and 9

contained 7 spots with Rf of 0.125 (blue), 0.3125 (red), 0.4375 (red), 0.5625 (purple), 0.6875 (red), 0.8125 (red), and 0.875 (blue), respectively. Similar to purified extract D, the purified extract E was produced from two fractions with six spots. The Rf values of each spot were 0.125 (blue), 0.3125 (red), 0.4375 (red), 0.5625 (purple), 0.6875 (red), and 0.875 (blue). Afterward, the steroid/triterpenoids compound of each purified extract was identified with Liebermann-Burchard reagent. This method was used to detect and analyze compounds in the sample [45]. Liebermann-Burchard reagent for detection of steroid and triterpenoid compounds, with positive tests for the appearance of green-blue for steroid and the appearance of red, brown, or purple for triterpenoid [46]. At the chromatogram, each purified extract had an appearance of green, brown, and purple which indicates the presence of steroid and triterpenoid compounds (Figure 2).




FIGURE 1 Thin layer chromatography (TLC) profile with gradient mobile phase (n-hexane: ethyl acetate) in silica gel F254 after being sprayed with 50% sulfuric acid

TABLE 3 The Retention factor (Rf) values of chromatograms after purification using vacuum liquid chromatography

Fraction Composition	weight (g)	Yield (%)	Many stains	Rf	Stain color	Purified extracts
Fraction 1	0	0	-	-	-	
Fraction 2	0.0537	1.79	2	0.8125; 0.875	Blue; Blue	A
Fraction 3	0.16	5.33	3	0.6875; 0.8125; 0.875	Red; Red; Red	B
Fraction 4	0.2316	7.72	6	0.3125; 0.4375; 0.5625; 0.6875; 0.8125; 0.875	Red; Red; Red purple; Red; Red; Blue	C
Fraction 5	0.2317	7.72	6	0.3125; 0.4375; 0.5625; 0.6875; 0.8125; 0.875	Red; Red; Red purple; Red; Red; Blue	
Fraction 6	0.1944	6.48	7	0.125; 0.3125; 0.4375; 0.5625; 0.6875; 0.8125; 0.875	Blue; Red; Red; Red purple; Red; Red; Blue	
Fraction 7	0.1523	5.07	7	0.125; 0.3125; 0.4375; 0.5625; 0.6875; 0.8125; 0.875	Blue; Red; Red; Red purple; Red; Red; Blue	
Fraction 8	0.1193	3.97	7	0.125; 0.3125; 0.4375; 0.5625; 0.6875; 0.8125; 0.875	Blue; Red; Red; Red purple; Red; Red; Blue	D
Fraction 9	0.0927	3.09	7	0.125; 0.3125; 0.4375; 0.5625; 0.6875; 0.8125; 0.875	Blue; Red; Red; Red purple; Red; Red; Blue	
Fraction 10	0.0732	2.44	7	0.125; 0.3125; 0.4375; 0.5625; 0.6875; 0.8125; 0.875	Blue; Red; Red; Red purple; Red; Red; Blue	
Fraction 11	0.2103	7.01	6	0.125; 0.1875; 0.3125; 0.375; 0.6875; 0.75	Blue; Red; Red; Red purple; Red; Red; Blue	E




FIGURE 2 Detection of steroids/triterpenoids compound of purified extract using thin layer chromatography (TLC) with Liebermann-Bourchard

Cytotoxic activity test from crude and purified extracts of Duku leaf was carried out against MCF-7 and HTB-183 cells using the MTT method. The absorbance was measured at 595 nm using a microplate reader. Absorbance obtained from the test results is used to calculate cell viability. The cancer cells were treated with various concentrations of samples ranging from 31.25 $\mu\text{g}/\text{mL}$ to 500.00 $\mu\text{g}/\text{mL}$. This study showed that the higher concentration gave the smaller % viability in living cancer cells (Figure 3). The 500.00 $\mu\text{g}/\text{mL}$ of samples have the strongest activity to kill cancer cells, even the % viability of MCF-7 and HTB-183 cells after treatment with the crude, purified extracts C, D, and E in this concentration is $0.00\% \pm 0.00$, respectively. After the concentration was decreased to 125.00 $\mu\text{g}/\text{mL}$, the purified extract D had a superior activity compared to the other samples in inhibiting the MCF-7 and HTB-183 cells of $0.00\% \pm 0.00$ and $4.67\% \pm 1.89$ with p

< 0.05 , respectively (Figure 3A and 3B). The inhibitory concentration 50 (IC_{50}) of each sample supported the purified extract D as the strongest sample than the other of $56.26 \pm 3.11 \mu\text{g}/\text{mL}$ against MCF-7 cells and $70.94 \pm 2.92 \mu\text{g}/\text{mL}$ against HTB-183 cells (Table 4). The lowest IC_{50} showed the best cytotoxic activity [47]. This phenomenon is not applied in all purified extracts. The IC_{50} showed that purified extracts have different cytotoxic activity against MCF-7 and HTB-183 cell lines. In these cases, only purified extract D has an IC_{50} value under 100 $\mu\text{g}/\text{mL}$ against two both cancer cells. Other purified extract was identified as similar to crude extract with IC_{50} in a range of 100-200 $\mu\text{g}/\text{mL}$. However, this study showed the purified extract A has IC_{50} more than 200 $\mu\text{g}/\text{mL}$ against MCF-7 and HTB-183 cell lines. This situation illustrated that the active compound from Duku leaf was contained in purified extract D more than other purified extract even the crude extract.




FIGURE 3 The % viability of MCF-7 and HTB-183 cells after treatment using crude and purified extracts for 24 hours. All data are described in mean \pm SD, $n = 3$. A: % viability of MCF-7 cells, B: % viability of HTB-183 cells

This study described the cytotoxic activity of crude and purified extracts against MCF-7 and HTB-183 cells, and the result is similar to the previous study. The ethanolic extract of Duku leaf was reported to have cytotoxic activity against HepG₂ with IC₅₀ of 19.93 ± 0.93 $\mu\text{g/mL}$. That was reported the extract has a mechanism to inhibit PI3K/Akt/mTOR pathways [29]. On the other hand, the ethanolic extract of Duku leaf identified a strong agent to inhibit the PANC-1 cells with IC₅₀ of 28.61 ± 0.13 $\mu\text{g/mL}$ [30]. One of the secondary metabolites that exerts anticancer activity is the terpenes group [48]. The mechanism of action of terpenoids as anticancer is to inhibit the cell cycle [49], inhibit cell proliferation [50], induce apoptosis [51], block the transition of epithelial cells to mesenchymal cells [52], inhibit migration and invasion of cancer cells [53], and modulate the immune system PD-L1 [54].

The crude and purified extracts showed antioxidant properties. Several concentrations were created from the crude

and purified extracts ranging from 31.25 $\mu\text{g/mL}$ to 500.00 $\mu\text{g/mL}$. The purified extract D showed the highest % scavenging DPPH activity compared to the other samples in 125.00 $\mu\text{g/mL}$ with the values of $72.99\% \pm 2.37$, significantly different with $p < 0.05$ (Figure 4). After IC₅₀ calculation, the purified extract D has stronger antioxidant properties than other samples of 53.65 ± 1.55 $\mu\text{g/mL}$, with $p < 0.05$ (Table 4). The antioxidant properties of purified extract D describe a kind of reason for the strongest cytotoxic activity between samples and prove a strong correlation between antioxidant properties and cytotoxic activity (Table 5). Many studies have illustrated the correlation of antioxidant properties with cytotoxic activity [55]. The presence of antioxidant compounds increases the activity to kill cancer cells [56]. Several studies show that antioxidant compounds like phenolic and flavonoid have a role in inducing some proteins to lead the cell cycle inhibition and apoptosis induction [57,58].




FIGURE 4 The % DPPH scavenging activity after treatment using crude and purified extracts. All data are described in mean \pm SD, n = 3.

TABLE 4 The inhibitory concentration 50 (IC₅₀) values of crude and purified extracts (All data described in mean \pm SD, n = 3)

Samples	IC ₅₀ (µg/mL)		DPPH Scavenging
	MCF-7	HTB-183	
Crude extract	114.56 \pm 2.80	142.68 \pm 9.97	110.24 \pm 1.80
Purified extract A	229.14 \pm 4.48	246.96 \pm 10.43	126.34 \pm 1.35
Purified extract B	123.44 \pm 6.74	142.77 \pm 8.91	117.87 \pm 2.05
Purified extract C	101.58 \pm 3.70	133.45 \pm 6.32	112.46 \pm 2.21
Purified extract D	56.26 \pm 3.11	70.94 \pm 2.92	53.65 \pm 1.55
Purified extract E	112.23 \pm 11.95	141.66 \pm 7.46	151.23 \pm 1.67

TABLE 5 Pearson's correlation and significant values between cytotoxic and antioxidant activities

	Antioxidant properties	Cytotoxic activity (MCF-7)	Cytotoxic activity (HTB-183)
Antioxidant properties	-	R = 0.986 <i>p</i> < 0.05	R = 0.952 <i>p</i> < 0.05
Cytotoxic activity (MCF-7)	R = 0.986 <i>p</i> < 0.05	-	-
Cytotoxic activity (HTB-183)	R = 0.952 <i>p</i> < 0.05	-	-

Conclusion

The present study described the cytotoxic activity of crude and purified extracts from Duku leaf against MCF-7 and HTB-183 cell lines. Based on the findings, the purified extract D has better cytotoxic and antioxidant activities than the crude extract and other purified extracts. These results described the opportunity to develop anti-breast and lung cancer agents. Further investigations will be needed to describe the active compounds in

purified extract D and provide an explanation or scientific reason for the anti-cancer mechanism of purified extract D through several tests such as cell cycle arrest inhibition, apoptosis induction, and protein expression test.

Acknowledgments

Universitas Sumatera Utara provided financial support for this research through the

TALENTA scheme 2022 (Grant No. 12/UN5.2.3.1/PPM/KP-TALENTA/2022).

Conflict of interest

All authors declare to have no conflict of interest.

Orcid:

Muhammad Fauzan Lubis:

<https://orcid.org/0000-0001-9651-904X>

Sumaiyah Sumaiyah:

<https://orcid.org/0000-0002-0702-4801>

Embun Suci Nasution:

<https://orcid.org/0000-0003-4763-6586>

Ririn Astyka:

<https://orcid.org/0000-0002-1832-5132>

Masfria Masfria:

<https://orcid.org/0000-0001-6087-4349>

Hafid Syahputra:

<https://orcid.org/0000-0001-6628-7727>

Carlos Nofanolo Lase:

<https://orcid.org/0009-0009-1561-1924>

References

- [1] J. Ferlay, M. Colombet, I. Soerjomataram, D.M. Parkin, M. Pineros, A. Znaor, F. Bray, Cancer statistics for the year 2020: An overview, *Int. J. Cancer*, **2021**, *149*, 778-789. [Crossref], [Google Scholar], [Publisher]
- [2] R.L. Siegel, K.D. Miller, N.S. Wagle, A. Jemal, Cancer statistics, 2023, *CA Cancer J Clin*, **2022**, *73*, 17-48. [Crossref], [Google Scholar], [Publisher]
- [3] A. Leiter, R.R. Veluswamy, J.P. Wisnivesky, The global burden of lung cancer: current status and future trends, *Nat Rev Clin Oncol*, **2023**, *20*, 624-639. [Crossref], [Google Scholar], [Publisher]
- [4] W. Ju, R. Zheng, S. Zhang, H. Zeng, K. Sun, S. Wang, R. Chen, L. Li, W. Wei, J. He, Cancer statistics in Chinese older people, 2022: current burden, time trends, and comparisons with the US, Japan, and the Republic of Korea, *China Life Sci*, **2023**, *66*, 1079-1091. [Crossref], [Google Scholar], [Publisher]
- [5] B. Ng, H. Puspitaningtyas, J.A. Wiranata, S.H. Hutajulu, I. Widodo, N. Anggorowati, G.Y. Sanjaya, L. Lazuardi, P. Sripan, Breast cancer incidence in Yogyakarta, Indonesia from 2008–2019: A cross-sectional study using trend analysis and geographical information system, *PloS one*, **2023**, *18*, e0288073. [Crossref], [Google Scholar], [Publisher]
- [6] O.D. Asmara, E.D. Tenda, G. Singh, C.W. Pitoyo, C.M. Rumende, W. Rajabto, N.R. Ananda, I. Trisnawati, E. Budiyo, H.F. Thahadian, E.C. Boerma, A. Faisal, D. Hutagaol, W. Soeharto, F. Radityamurti, E. Marfiani, P.Z. Romadhon, F.N. Kholis, H. Suryadinata, A.Y. Soeroto, S.A. Gondhowiardjo, W.H. van Geffen, Lung cancer in Indonesia, *J. Thorac Oncol.*, **2023**, *18*, 1134-1145. [Crossref], [Google Scholar], [Publisher]
- [7] S. Hashem, T.A. Ali, S. Akhtar, S. Nisar, G. Sageena, S. Ali, S. Al-Mannai, L. Therachiyil, R. Mir, I. Elfaki, M.M. Mir, F. Jamal, T. Masoodi, S. Uddin, M. Singh, M. Haris, M. Macha, A.A. Bhat, Targeting cancer signaling pathways by natural products: Exploring promising anti-cancer agents, *Biomed. Pharmacother.*, **2022**, *150*, 113054. [Crossref], [Google Scholar], [Publisher]
- [8] S.M. Kasim, N.T. Abdulaziz, Y.F. Mustafa, Synthesis and biomedical activities of coumarins derived from natural phenolic acids, *J. Med. Chem. Sci.*, **2022**, *5*, 546-560. [Crossref], [Google Scholar], [Publisher]
- [9] Y.F. Mustafa, M.K. Bashir, M.K. Oglah, Influence of albobcarbon-cylic hybridization on biomedical activities: a review, *J. Med. Chem. Sci.*, **2022**, *5*, 550-568. [Crossref], [Publisher]
- [10] Elfahmi, H.J. Woerdenbag, O. Kayser, Jamu: Indonesian traditional herbal medicine towards rational phytopharmacological use, *J. Herb. Med.*, **2014**, *4*, 51-73. [Crossref], [Google Scholar], [Publisher]
- [11] A. Oladipupo, C. Alaribe, T. Akintemi, H. Coker, Effect of *Phaulopsis falcisepala* (Acanthaceae) leaves and stems on mitotic arrest and induction of chromosomal changes in Meristematic Cells of *Allium Cepa*, *Prog*

- Chem Biochem Res*, **2021**, *4*, 134-147. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [12] C. Ukwubile, E. Ikpefan, M. Bingari, L. Tam, Acute and subchronic toxicity profiles of *Melastomastrum capitatum* (Vahl) Fern. (Melastomataceae) root aqueous extract in Swiss albino mice, *Prog Chem Biochem Res*, **2019**, *2*, 74-83. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [13] P.A. Kalvanagh, Y.A.A. Kalvanagh, New therapeutic approach based on silymarin in the treatment of breast cancer, *Adv. J. Chem. Sect. B. Nat. Prod. Med. Chem.*, **2023**, *5*, 75-85. [[Crossref](#)], [[Pdf](#)], [[Publisher](#)]
- [14] L. Ghaderi, Relationship between FGFR2 gene RS2981582 polymorphism and breast cancer risk factors in women candidates for surgery, *Adv. J. Chem. Sect. B. Nat. Prod. Med. Chem.*, **2023**, *5*, 98-107. [[Crossref](#)], [[Pdf](#)], [[Publisher](#)]
- [15] M. Zehravi, M. Maqbool, I. Ara, Curcumin—A promising phytochemical of immense potential, *Adv. J. Chem. Sect. B. Nat. Prod. Med. Chem.*, **2021**, *3*, 271-276. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [16] M. Greenwell, P.K. Rahman, Medicinal plants: their use in anticancer treatment, *Int J Pharm Sci Res*, **2015**, *1*, 4103-4112. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [17] F.Z. Kazemabadi, A. Heydarinasab, A. Akbarzadehkhayavi, M. Ardjmand, Development, optimization and in vitro evaluation of etoposide loaded lipid polymer hybrid nanoparticles for controlled drug delivery on lung cancer, *Chem. Methodol.*, **2020**, *5*, 135-152. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [18] M.F. Lubis, P.A.Z. Hasibuan, H. Syahputra, R. Astyka, A review on phytochemicals and pharmacological activities as ethnomedicinal uses of Duku (*Lansium domesticum* Corr.), *Open Access Maced J Med Sci*, **2022**, *10*, 57-65. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [19] A. Manosroi, P. Jantrawut, M. Sainakham, W. Manosroi, J. Manosroi, Anticancer activities of the extract from Longkong (*Lansium domesticum*) young fruits, *Pharm Biol*, **2012**, *50*, 1397-1407. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [20] S.E. Sinaga, F.F. Abdullah, U. Supratman, T. Mayanti, R. Maharani, Isolation and Structure Determination of Stigmaterol from the Stem Bark of *Lansium domesticum* Corr. Cv. Kokossan, *Chimica et Natura Acta*, **2022**, *10*, 106-111. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [21] R.M.A. Khalili, J.M. Noratiqah, R. Norhaslinda, A.H. Norhayati, B.A. Amin, A. Roslan, A.L.A. Zubaidi, Cytotoxicity effect and morphological study of different Duku (*Lansium domesticum* corr.) extract towards human colorectal adenocarcinoma cells line (HT-29), *Pharmacogn. J.*, **2017**, *9*, 757-761. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [22] H.M. Abdallah, G.A. Mohamed, S.R.M. Ibrahim, *Lansium domesticum*—a fruit with multi-benefits: traditional uses, phytochemicals, nutritional value, and bioactivities, *Nutrients*, **2022**, *14*, 1531. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [23] A.V. Lee, S. Oesterreich, N.E. Davidson, MCF-7 cells changing the course of breast cancer research and care for 45 years, *J. Natl. Cancer Inst.*, **2015**, *107*, 1-4. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [24] A. Moghaddam, H.A. Zamani, H. Karimi-Maleh, A new sensing strategy for determination of tamoxifen using Fe₃O₄/graphene-ionic liquid nanocomposite amplified paste electrode, *Chem Methodol*, **2021**, *5*, 373-380. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [25] A. Maatta, K. Makinen, A. Ketola, T. Liimatainen, F.N. Yongabi, M. Vaha-Koskela, R. Pirinen, O. Rausti, R. Pellinen, A. Hinkkanen, J. Wahlfors, Replication competent semliki forest virus prolongs survival in experimental lung cancer, *Int. J. Cancer*, **2008**, *123*, 1704-1711. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [26] A. Amann, M. Zwierzina, G. Gamerith, M. Bitsche, J.M. Huber, G.F. Vogel, M. Blumer, S. Koeck, E.J. Pechriggl, J.M. Kelm, W. Hilbe, H. Zwierzina, Development of an innovative 3D cell culture system to study tumour – stroma

- interactions in non-small cell lung cancer cells, *Plosone*, **2014**, *9*, e92511. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [27] K. Fadhilah, S. Wahyuono, P. Astuti, A bioactive compound isolated from Duku (*Lansium domesticum* Corr) fruit peels exhibits cytotoxicity against T47D cell line, *F1000Res*, **2020**, *9*, 3. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [28] K. Fadhilah, S. Wahyuono, P. Astuti, Fractions and isolated compounds from *Lansium domesticum* fruit peel exhibited cytotoxic activity against T-47D and HepG2 cell lines, *Biodeversitas*, **2021**, *22*, 3743-3748. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [29] M.F. Lubis, P.A.Z. Hasibuan, H. Syahputra, J.M. Keliat, V.E. Kaban, R. Astyka, Duku (*Lansium domesticum*) leaves extract induces cell cycle arrest and apoptosis of HepG2 cells via PI3K/Akt pathways, *Trends in Sciences*, **2022**, *20*, 6437. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [30] M.F. Lubis, P.A.Z. Hasibuan, V.E. Kaban, R. Astyka, Phytochemicals analysis And cytotoxic activity of *Lansium Domesticum* corr extract-cisplatin combination against panc-1 cell line, *Rasayan J Chem*, **2023**, *16*, 32-37. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [30] M.F. Lubis, V.E. Kaban, J.O. Aritonang, D. Satria, A.A. Mulina, H. Febriani, Acute toxicity and antifungal activity of the ointment *Murraya koenigii* ethanol extract, *Rasayan J. Chem*, **2022**, *15*, 256-261. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [31] M.S. Fareza, N.A. Choironi, S.S. Susilowati, M.P. Rini, V. Festihawa, I.S.N. Fauzi, E.D. Utami, Sarmoko, LC-MS/MS analysis and cytotoxic activity of extract and fractions of *Calophyllum soulattri* stembark, *Indonesian J Pharm*, **2021**, *32*, 356-364. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [32] K.S. Sasmitaloka, S.M. Widayanti, I. Mulywanti, E.S. Iriani, Physicochemical and antioxidant characteristics of black garlic from indigenous Indonesian garlic, *IOP Conf. Ser.: Earth Environ. Sci*, **2022**, *1041*, 012004. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [33] C.K. Jiea, S. Fuloria, V. Subrimanyan, M. Sekar, K.V. Sathasivam, S. Kayarohanam, Y.S. Wu, V.S.S.R. Velaga, A.K. Janakiraman, M.N.H. Maziz, N.K. Fuloria, Phytochemical screening and antioxidant activity of *Cananga odorata* extract, **2022**, *15*, 1230-4. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [34] P.A.Z. Hasibuan, M.F. Lubis, J.M. Keliat, Cytotoxic test combination of ethyl acetate extract african leaves (*Vernonia amygdalina* Delile) and gemcitabine on PANC-1 cells, *AIP Conf. Proc.* **2023**, *2626*, 030004. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [35] K. Fitri, M. Andry, T.N. Khairani, H.S. Winata, A. Violenta, N. Lubis, M.F. Lubis, Synthesis of silver nanoparticles using ethanolic extract of *Nelumbo Nucifera* Gaertn. leaf and its cytotoxic activity against T47D and 4T1 cell lines, *Rasayan J. Chem*, **2023**, *16*, 104-110. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [36] M.F. Lubis, H. Syahputra, D.N. Illian, V.E. Kaban, Antioxidant activity and nephroprotective effect of *Lansium parasiticum* leaves in doxorubicin-induced rats, *J Res Pharm*, **2022**, *26*, 565-573. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [37] I.P. Sany, Romadhon, A.S. Fahmi, Physicochemical characteristics and antioxidant activity of solid soap enriched with crude *Eucheuma cottoni* Extract, *IOP Conf Ser: Earth Environ Sci*, **2018**, *246*, 012066. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [38] L. Nurhidayati, Y. Fitriani, S. Abdillah, E. Mumpuni, M. Rafi, Physicochemical properties and antioxidant activities of crude fucoidan extracted from *Sargassum cinereum*, *Indones. J. Pharm. Sci.*, **2020**, *18*, 68-74. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [39] T.S. Wahyuni, N. Khoiriyah, L. Tumewu, W. Ekasari, A. Fuad, A. Widyawaruyanti, Microscopic and physicochemical evaluation of *Ruta angustifolia* leaves, *J. Public Health Afr.*, **2023**, *14*, 2520. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [40] V.L. Thi, N. Nguyen, Q. Nguyen, Q.V. Dong, T. Do, K.T. Nguyen, Phytochemical screening and potential antibacterial activity of defatted

- and nondefatted methanolic extracts of Xao Tam Phan (*Paramignya trimera* (Oliv.) Guillam) peels against multidrug-resistant bacteria, *Scientifica (Cairo)*, **2021**, 34513112. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [41] R. Yuniati, M. Zainuri, H. Kusumaningrum, Qualitative tests of secondary metabolite compounds in ethanol extract of *Spirulina platensis* from Karimun Jawa sea, Indonesia, *Biosaintifika*, **2020**, *12*, 343-349. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [42] R. Purnama, A. Primadhamanti, Phytochemical screening, spectrum profile of functional groups, and effervescent formulation of kepok banana peels stem extract, *ALKIMIA: Jurnal Ilmu Kimia dan Terapan*, **2021**, *4*, 66-72. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [43] P. Praptiwi, D. Wulansari, A. Fathoni, N. Harnoto, R. Novita, Alfridsyah, A. Agusta, Phytochemical screening, antibacterial and antioxidant assessment of *Leuconotis eugenifolia* leaf extract, *Nusantara Bioscience*, **2020**, *12*, 79-85. [[Crossref](#)], [[Google Scholar](#)]
- [44] A.G. Fasya, S. Amalia, D.S. Megawati, V.A. Kusuma, B. Purwantoro, Isolation, identification, and bioactivity of steroids isolates from *Hydrilla verticillata* petroleum ether fraction, *IOP Conf Series: Earth and Environmental Science*, **2020**, *456*, 012009. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [45] S.A. Bhawano, O. Sulaiman, R. Hashim, M.N.M. Ibrahim, Thin-layer chromatographic analysis of steroids: a review, *Trop J Pharm Res*, **2010**, *9*, 301-313. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [46] M.F. Lubis, P.A.Z. Hasibuan, U. Harahap, D. Satria, H. Syahputra, M. Muhammad, R. Astyka, The molecular approach of natural products as pancreatic cancer treatment: a review, *Rasayan J. Chem.*, **2022**, *15*, 1362-1377. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [47] S. Kamran, A. Sinniah, M.A.M. Abdulgani, M.A. Alshawsh, Therapeutic potential of certain terpenoids as anticancer agents: a scoping review, *Cancers (basel)*, **2022**, *14*, 1100. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [48] H. Chen, K. Lin, A. Huang, H. Tu, B. Wei, T. Hour, M. Yen, Y. Pu, C. Lin, Terpenoids induce cell cycle arrest and apoptosis from the stems of *Celastrus kusanoi* associated with reactive oxygen species, *J Agric Food Chem*, **2010**, *58*, 3808-3812. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [49] Y. Liu, R.J. Whelan, B.R. Pattnaik, K. Ludwig, E. Subudhi, H. Rowland, N. Claussen, N. Zucker, S. Uppal, D.M. Kushner, M. Felder, M.S. Patankar, A. Kapur, Terpenoids from *Zingiber officinale* (Ginger) induce apoptosis in endometrial cancer cells through the activation of p53, *Plos One*, **2012**, *7*, e53178. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [50] R. Mutiah, A. Widyawaruyanti, S. Sukardiman, Calotroposid A: a Glycosides Terpenoids from *Calotropis gigantea* induces apoptosis of colon cancer WiDr cells through cell cycle arrest G2/M and caspase 8 expression, *Asian Pac. J. Cancer Prev.*, **2018**, *19*, 1457-1464. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [51] J. Fu, S. Wang, H. Lu, J. Ma, X. Ke, T. Liu, Y. Luo, *In vitro* inhibitory effects of terpenoids from *Chloranthus multistachys* on epithelial-mesenchymal transition via down-regulation of Runx2 activation in human breast cancer, *Phytomedicine*, **2015**, *22*, 165-172. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [52] H. Sun, L. Zhang, B. Sui, Y. Lo, J. Yan, P. Wang, Y. Wang, S. Liu, The effect of terpenoid natural chinese medicine molecular compound on lung cancer treatment, *Evid. Based Complementary Altern. Med.*, **2021**, 3730963. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [53] C. Bailly, G. Vergoten, Proposed mechanisms for the extracellular release of PD-L1 by the anticancer saponin platycodin D, *Int. Immunopharmacol.*, **2020**, *85*, 106675. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [54] M. Rajan, G. Rajkumar, T.J.F.L. Guedes, R.G.C. Barros, N. Narain, Performance of different solvents on extraction of bioactive compounds, antioxidant and cytotoxic

activities in *Phoenix loureiroi* Kunth leaves, *J. Appl. Res. Med. Aromat. Plants*, **2020**, *17*, 100247. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]

[55] N. Armania, L.S. Yazan, S.N. Musa, I.S. Ismail, J.B. Foo, K.W. Chan, H. Noreen, A.H. Hisyam, S. Zulfahmi, M. Ismail, *Dillenia suffruticosa* exhibited antioxidant and cytotoxic activity through induction of apoptosis and G₂/M cell cycle arrest, *J. Ethnopharmacol.*, **2013**, *146*, 525–535. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]

[56] X. Chen, J. Shen, J. Xu, T. Herald, D. Smolensky, R. Perumal, W. Wang, Sorghum phenolic compounds are associated with cell growth inhibition through cell cycle arrest and apoptosis in human hepatocarcinoma and colorectal adenocarcinoma cells, *Foods*, **2021**, *10*, 993. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]

[57] P.F. Rezaei, S. Fouladdel, S. Hassani, F. Yousefbeyk, S.M. Ghaffari, G. Amin, E. Azizi, Induction of apoptosis and cell cycle arrest by pericarp polyphenol-rich extract of *Baneh* in human colon carcinoma HT29 cells, *Food Chem. Toxicol.*, **2012**, *50*, 3-4. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]

How to cite this article: Muhammad Fauzan Lubis*, Sumaiyah Sumaiyah, Embun Suci Nasution, Ririn Astyka, Masfria Masfria, Hafid Syahputra, Carlos Nofanolo Lase. Cytotoxic activity of the purified extracts from duku (*Lansium domesticum* Corr.) Leaf against MCF-7 and HTB-183 cell lines, and the correlation with antioxidant activity. *Eurasian Chemical Communications*, 2023, 5(12), 1082-1095. **Link:** https://www.echemcom.com/article_182179.html