

FULL PAPER

The role of oxidative stress in hypoalbuminemia nephropathy related to Nephrotic syndrome: a critical review

Amallia Nuggetsiana Setyawati^{a,b,c} 

^aDepartment of Medical Biology and Biochemistry, Faculty of Medicine Universitas Diponegoro, Semarang Indonesia Semarang-50275, Central Java, Indonesia

^bDepartment of Pediatrics, Diponegoro National Hospital, Semarang-50275, Central Java, Indonesia

^cDivision of Pediatrics, Williambooth General Hospital, Semarang-50131, Central Java, Indonesia

Nephrotic syndrome (NS) is defined as severe proteinuria that results in low albumin levels, increased permeability within the glomerular filtration barrier, and functional impairment. Oxidative stress and inflammation are believed to play a significant role in the development of nephropathy. Reactive oxygen species, mitochondria, nitric oxide (NO) synthases, and xanthine oxidases are all injured by kidney-inducing substances. This study aimed to elucidate the mechanism of hypoalbuminaemia in nephrotic syndrome related to oxidative stress in NS. Through studies and reviews, the causes, pathophysiology, sources, and agents of renal oxidative stress have been elucidated over several decades. We reviewed studies on reactive oxygen species (ROS) formation and their relationship with hypoalbuminaemia nephrotic syndrome. The pathogenic pathways that lead to renal fibrosis, mechanisms of oxidative stress production during renal disorders, and medications that specifically target oxidative stress during tubulointerstitial fibrosis and glomerulosclerosis are explained in this article. A distinguishing feature of NS is increased excretion of albumin and other serum proteins. Therapies that target oxidative stress have the potential to treat renal fibrosis, given the importance of oxidative stress in renal nephrotic syndrome.

***Corresponding Author:**

Amallia Nuggetsiana Setyawati
Email: amallia.setyawati@fk.undip.ac.id
Tel.: +081326294457

KEYWORDS

Hypoalbuminemia; nephrotic syndrome; oxidative stress.

Introduction

Nephrotic syndrome (NS) is defined as severe proteinuria that results in low albumin levels, increased permeability within the glomerular filtration barrier, and functional impairment. Although glucocorticoids are the cornerstone of NS processing, they cause significant side effects and resistance in many patients, thereby increasing the risk of chronic or end-stage renal impairment. The aetiology of NS is complex and includes immunological dysfunction, oxidative stress, and molecular changes in the podocytes. Free radicals in the

human body are known oxidants. Oxidants in the human body include endogenous and/or exogenous oxidants. The human body functions as an oxidant. These ants include superoxide anions (O_2^-), reactive oxygen species (ROS), hydroxyl radicals (OH^-), singlet oxygen (1O_2), and hydrogen peroxide (H_2O_2) [1]. Oxidants in the human body include endogenous and/or exogenous oxidants [2].

Free radicals and other ROS can be produced by both internal and external factors such as exposure to the use of X the ozone layer, cigarette smoke, air pollution, and industrial toxic substances. Free radicals are

continuously produced within cells via enzymatic and nonenzymatic activities. Several enzymatic processes can produce free radicals, including the cytochrome P-450 framework, prostaglandin phagocytosis, and respiratory chains [3]. Prooxidants can also be produced by ionising or nonenzymatic interactions between oxygen and organic molecules. All molecules of oxygen (O_2) are included in the word "prooxidants", and they have a higher reactivity than superoksida, hydroxyl ($HO\cdot$), and hydrogen peroxide (H_2O_2). ROS is produced in the tubules of the body via normal aerobic metabolism. Oxidants' nature was somewhat more stable than that of non-oxidant compounds, but with significantly higher levels of activity [4]. This review article tries to figure out how oxidative stress is linked to hypoalbuminemic nephropathy and the progress made in making antioxidants that can lessen the damage to podocytes that happens as a result. The endothelium, podocytes, and foot processes line glomerular capillaries, forming a glomerular filter. The larger proteins were eliminated after filtering. Podocyte damage can result in irreversible damage to the glomerulus. Proteinuria can occur because of injury to the basement membrane of the glomerular system, epithelial surface, or podocytes. Albumin, the major component, accounts for 85% of proteinuria cases. Nonselective proteinuria induced by glomerular permeability abnormalities could not be distinguished. Therefore, this review aimed to determine how oxidative stress affects hypoalbuminaemia and nephropathy related to nephrotic syndrome.

Nephrotic syndrome, clinical spectrum, and laboratory hallmarks

Children aged > 1 year with NS (>80%) and steroid-sensitive nephrotic syndrome (NSSS) in remission after steroid therapy often show full enhancement in accordance with symptoms. However, steroid-resistant

nephrotic syndrome (NSRS) is found in 20% of children older than one year, and the majority of infants are younger than one year. However, some children with steroid-sensitive nephrotic syndrome (NSSS) experience remission [5,6].

Many proteins in the urine (40 mg/m² LPB/h, urine protein/creatinine ratio > 2 mg/mg, or dipstick 2+) are signs of nephrotic syndrome (NS), a medical condition. Other symptoms include oedema, hypoalbuminemia (2.5 g/dL), and the potential presence of hypercholesterolaemia [7]. High homocysteine levels and insufficient vitamin B levels can lead to thrombosis. NS is a condition that requires strict parental home oversight, adjustment of medication dosages, and daily intake protocols, which can lead to relapse and increased complexity. Our previous study indicated that the intervention group had somewhat higher average medication adherence compliance [8]. Children who lose protein in their urine may develop life-threatening consequences such as infection and thrombosis. Within the first three months of life, congenital nephrotic syndrome develops, infantile nephrotic syndrome develops between four and 12 months of life, and paediatric nephrotic syndrome develops beyond one year of life [9]. The clinical spectrum of NS is shown in Figure 1.

Childhood NS is divided into three categories: congenital (less than 1%), secondary (10%), and idiopathic (INS; 90% of cases). The two primary histological forms of idiopathic NS were MCD (85%) and FSGS (Focal Segmental Glomerulosclerosis (FSGS) (10%). Systemic conditions, including Alport syndrome, Henoch-Schonlein purpura nephritis, and lupus nephritis, are considered secondary NS. Severe proteinuria before the age of three months is referred to as congenital NS and is associated with CMV, syphilis, or other congenital infections as well as abnormalities in the genes that code for podocyte proteins (NPHS1: NPHS1: Congenital Nephrotic Syndrome of the Finnish

type, NPHS2: NPHS2 Stomatin Family Member, and WT1: Wilms tumour protein gene) [5,10].




FIGURE 1 Clinical signs and hallmarks of Nephrotic syndrome (NS)

Hypoalbuminemia in Nephrotic syndrome

Increased glomerular capillary wall permeability, which results in severe proteinuria and hypoalbuminaemia, is the primary anomaly observed in NS patients. Because of the extreme flattening of podocyte foot processes (a characteristic of idiopathic nephrotic syndrome) observed in patients with NS, podocytes may be significant in the biopsies of these patients [10,11]. Complex immune system diseases, particularly T cell-mediated immunity, have been linked to idiopathic nephrotic syndrome. A subgroup of activated lymphocytes creates plasma factors that increase capillary wall permeability in focal segmental glomerulosclerosis (FSGS). Podocyte protein (podocin and -actinin 4) and podocyte gene MYH9 mutations are also linked to focal segmental glomerulosclerosis (FSGS) [12].

In nephrotic syndrome, the complex interaction between free fatty acids (FFAs) and albumin changes the molecular link between proteinuria and high triglyceride levels. FFAs are mostly used as fuel by skeletal muscles and the heart. These organs take in both albumin-bound FFAs from food and lipolysis, and FFAs are released by lipoprotein

lipase (LPL)-mediated hydrolysis. When low-FFA-content albumin is lost through urine, it is replaced by high-FFA-content albumin, which changes the ratio of FFA uptake. This causes the localised overexpression of Angptl4, which inactivates LPL in tissues and reduces FFA production and hypertriglyceridaemia. The rapid absorption of albumin-bound FFA in peripheral organs and the disproportionate retention of albumin at high FFA concentrations are required for Angptl4's antiproteinuric and hypertriglyceridemia-inducing actions. The NPHS2 (podocin) and WT1 genes, as well as other parts of the glomerular filtration system, such as the slit pore containing nephrin, NEPH1, and CD-2-related proteins, can be mutated in steroid-resistant nephrotic syndrome. Hypoalbuminemia Hypoalbuminemia is the systemic condition that proteinuria is most closely associated with. If the albumin level is less than 2.5 g/dL, hypoalbuminaemia is one of the signs and symptoms of nephrotic syndrome in children [13]. The liver usually produces 12-14 g of albumin per day (130-200 mg/kg), which is metabolised by the same metabolite. After filtered albumin is absorbed, 10% is catabolised in the proximal tubule of the kidney, where it occurs extrarenally [14].

Hypoalbuminaemia is a symptom of enhanced albumin catabolism and excessive urine protein loss in patients with nephrotic syndrome. Albumin loss through urine is a major factor in the occurrence of hypoalbuminaemia [15]. This is not the only cause of albumin loss through urination in individuals with nephrotic syndrome, although the rate of albumin production can increase by at least twofold. Hypoalbuminaemia is also assumed to be influenced by increased albumin loss in the gastrointestinal tract; however, there is scant evidence to support this claim [16]. Therefore, an adequate link between the decline in the rate of albumin synthesis in the liver and the subsequent increase in albumin catabolism is required for the development of hypoalbuminaemia. While studies on patients with hypoalbuminaemia related to nephrotic syndrome have demonstrated that the rate of albumin production in the liver is only slightly above normal, under normal circumstances, the rate of albumin synthesis in the liver may increase up to 300% [17], indicating that the liver does not respond adequately to albumin synthetically. The plasma oncotic pressure, which perfuses the liver, is a key regulator of protein production. A two-fold increase in the rate of hepatic albumin gene transcription compared with normal is experimental data suggesting a hereditary shortage of circulating albumin. However, the increase in albumin synthesis in the liver was insufficient to offset the degree of hypoalbuminaemia, indicating a compromised synthetic response [18].

This also occurs in patients with NS; a drop in oncotic pressure is inadequate to speed up liver albumin production enough to increase plasma albumin concentration. In addition, there is evidence that albumin synthesis is controlled by the hepatic interstitial albumin levels in healthy individuals. The response to albumin production is normal and increases slightly but insufficiently in nephrotic syndrome because the hepatic interstitial albumin pool is not depleted. Albumin

production is affected by dietary protein consumption [19] and an increase in hepatic protein and albumin mRNA synthesis, whereas a low-protein diet does not [20]. Hyperfiltration caused by increased protein ingestion also results in increased albuminuria, whereas the serum albumin levels are unaffected. There is an ongoing debate regarding whether renal albumin catabolism causes hypoalbuminaemia in nephrotic syndrome. An increase in filtered protein that is only released in the urine, rather than being taken in and digested, indicates that the ability of renal tubules to transport albumin has reached its saturation threshold at healthy levels of processed albumin [21]. Studies on the perfusion of isolated proximal tubules in rabbits have shown that albumin absorption uses a dual transport system [22]. In this study, there are two types of structures: a low-capacity system that stops working when there is too much protein, and a high-capacity system with low affinity that allows the tubular rate of albumin absorption to increase as the filtered load increases. This theory is reinforced by the positive relationship between albumin fraction catabolism and albuminuria caused by puromycin aminonucleoside PAN (Puromycin Aminonucleoside-Induced Nephrosis) [23,24]. However, the absolute catabolic rate may be normal or even lower in nephrotic syndrome because of significantly decreased total body albumin storage. The fact that absolute albumin catabolism is decreased in nephrotics on a low-protein diet, but not when they consume a regular amount of protein, is proof that this has an impact on their nutritional status [25]. It is now obvious that numerous modifications in albumin homeostasis that cause low albumin levels in nephrotic syndrome cannot be completely compensated for by decreasing tubular renal albumin catabolism and boosting hepatic albumin synthesis. Several theories have been proposed to explain the occurrence of oedema in nephrotic individuals. The subjects of this

study were the traditional hypothesis regarding oedema formation [26]. Accordingly, fluid seeps into the interstitial space because of oedema caused by a drop in the intravascular oncotic pressure. Albumin leakage and hypoalbuminaemia are caused by an increase in glomerular capillary permeability. The determination of oncotic pressure is an essential activity of albumin. Therefore, the intravascular plasma colloid oncotic pressure decreases owing to hypoalbuminaemia. Oedema occurs when transudate fluid from the intravascular area leaks past the capillary walls into the interstitial space [17]. Figure 2 depicts the pathophysiology and mechanism of oedema development in patients with NS. According to the overflow hypothesis, 16 primary intrarenal processes regulate renal salt and water retention, as opposed to peripheral systemic stimulation. Primary renal salt

retention causes a spike in the amount of extracellular fluid and the plasma. When there is too much fluid in the interstitial space, oedema develops [27]. Hypercholesterolaemia-nephropathy syndrome is characterised by an increase in almost all lipid levels (cholesterol, triglycerides, and serum lipoproteins). There are several possible explanations for this, including hypoproteinaemia, which increases the synthesis of all proteins in the liver, including lipoproteins [28]. A decrease in plasma levels of lipoprotein lipase, the primary enzyme system that removes fat from the plasma, also contributes to a decrease in fat catabolism. Hypoalbuminaemia, hyperlipidaemia, and peripheral oedema are caused by reduced glomerular capillary wall permeability, which also results in the failure to maintain protein loss in the urine [29].




FIGURE 2 Pathophysiology and oedema mechanism of nephrotic syndrome

Several clinical disorders have been associated with free radical oxidative damage, in kidney injury [30,31].

Stress oxidative role in Nephrotic syndrome




FIGURE 3 Oxidative stress and podocyte-related injuries alter the glomerular filtration barrier. ROS: reactive oxygen species, RNS: reactive nitrogen species

Glomerular illnesses with diverse aetiologies are characterised by albuminuria. It is a cause of glomerular disease and is not only one of its symptoms, but also triggers interstitial fibrosis, glomerulosclerosis, and eventually a loss in kidney function. Reactive oxygen species (ROS) affect glomerular haemodynamics, decrease tubular epithelial cell structural integrity, and increase glomerular responsiveness to proteins, all of which contribute to cell damage via lipid peroxidation. Antioxidants which protect podocytes from ROS-induced cell death can be used to treat podocytopathy, as this system is a major factor in the emergence of podocytopathies. As free radicals are oxidative stressors in a number of disorders, their impact on the cause of nephrotic syndrome (NS) is significant [32]. The basis of this study was oxidative damage caused by free radicals, which has been associated with an assortment of clinical illnesses, including kidney damage [33]. Figure 3 shows the interaction between oxidative stress and podocyte-related injury. Children with N.S. are more likely to

experience recurrence because they are more susceptible to repeated infections. Therefore, an individual's vulnerability to sickness may be influenced by reductions in antioxidants, including superoxide dismutase (SOD), glutathione (GSH), plasma (ceceruloplasmin (, and antioxidant vitamin E. These antioxidants also aid the immune system by reducing harm and adjusting to challenging circumstances [34,35]. Idiopathic nephrotic syndrome (NSI) is a T-cell disorder in which cells release cytokines that affect the glomeruli and improve plasma protein permeability. A particular type of T cell that aids in maintenance is known as a regulatory T cells [36]. Several illnesses are managed with T-cell function inhibitors such as cyclosporine and corticosteroids. Immune system failure can cause selective proteinuria, which reduces the amount of circulating substances that modify the diaphragm gap [37].

Focal segmental glomerulosclerosis (FSGS) and minimal change disease are two major symptoms of NS. MCD is the most common cause of NS among children. The exact cause of

NS remains unknown [38]. However, based on the response of NS to immunosuppressive drugs such as corticosteroids and calcineurin inhibitors, which modulate T cell function, NS is believed to result from primary T cell failure [39].

A redox protein called CHOP-TXNIP (C/EBP homologous protein)-TXNIP (thioredoxin-interacting protein) is crucial for both mitochondrial dyshomeostasis and ER failure, the study's findings indicate. NS

triggers mitochondrial oxidative stress (ROS). Through triggering TXNIP translocation from the nucleus to the mitochondria, CHOP up-regulates Trx2 and frees ASK1. CHOP removal inhibits TXNIP translocation and ROS overproduction, suggesting a possible therapeutic strategy for NS [40]. Table 1 summarises nephrotic syndrome supplementation that may be therapeutically useful for improving nephropathy and its proposed mechanism.

TABLE 1 Nephrotic syndrome supplementation that may be therapeutically useful

No.	Molecules	Animal experimentation	Treatment	Mechanism	Ref.
1.	Co-Q10 (CO-Q10)	There have been reports of B6/Pdss2kd/kd mutant mice. It is thought that these mice have a spontaneous mutation in the gene that codes for polyprenyl-diphosphate synthase (Pdss2) subunit 2.	CoQ10 supplementation of COQ8A depleted cells	Co-Q10 and other antioxidants protect podocytes from ROS-mediated cell death. These antioxidants could be used to treat people with podocytopathy because this process is important for understanding how the disease works.	[41,42]
2.	Co Q10	116 patients of SNRS patients	Q-10 supplementation was administered to the patients in two or three doses per day, with doses ranging from 3 to 60 mg/kg.	According to the study, using a CoQ10 supplement at 12 months dramatically decreased proteinuria by 88%.	[43]
3.	<i>Zingiber officinale</i>	Cisplatin induced nephrotoxicity animal models	Hydroalcoholic extract of <i>Stachys pilifera</i> Benth. (500 mg kg ⁻¹ day ⁻¹) was orally administered.	<i>Z. officinale</i> significantly and dose-dependently prevented cisplatin-induced nephrotoxicity.	[44]
4.	Zinc	Young idiopathic NS patients	Zinc sulphate (10 mg / day	A potential inclusion in steroid-sensitive nephrotic syndrome treatment regimens is suggested by the increased continuous remission rates of	[45]

5.	Combination of α -tocopherol and vitamin C	Young idiopathic NS patients	For a period of 12 weeks, oral vitamin E (D1- α -tocopherol acetate) and vitamin C (Sigma-Aldrich, USA) were given at doses of 10-15 mg/kg body weight (maximum dose 400 mg) and 10-15 mg/kg body weight, respectively.	zinc supplementation. When α -tocopherol and vitamin C were given together to young people with idiopathic nephrotic syndrome, their blood pressure went down in both prehypertensive and hypertensive people, but there was no significant change in NO _x and ox-LDL levels.	[46]
6.	Curcumin	Doxorubicyn induced models	Curcumin (oral gavage, once day, 50 mg/kg)	Curcumin increased Nrf2, reduced NF- κ B, and increased podocin. In addition, it relieved proteinuria, reduced blood lipid levels, and improved hypoalbuminemia in NS rats. Curcumin markedly improved renal function and podocyte destruction by lowering blood viscosity and hematocrit. It repaired kidney damage caused by oxidation by boosting the production of antioxidant enzymes and switching on Nrf2.	[47]
7.	Combination of <i>C.xanthorrhiza</i> Roxb and <i>Nigella sativa</i>	Adriamycin induced mouse models	Black cumin (250 mg/kg BW daily) and temulawak (1000 mg/kg BW daily) rhizome preparations	A study found that using <i>Nigella sativa</i> extract and <i>Curcuma xanthorrhiza</i> rhizome together lowered cholesterol and triglyceride levels in male Wistar rats. However, the NS model rats did not show any significant changes.	[48]

Many studies have attempted to determine how NS regulation affects the expression, operation, and release of cytokines on T-cell surfaces. Several other networks have been suggested to be connected to the NS. TNF, CLC-1, and IL-18 are the three cytokines that cause tissue damage [49]. The expression of sphingomyelin phosphodiesterase acid-like 3b (SMPDL3b), angiopoietin-like 4 (Angptl4), and glomerular podocyte B7-1 (CD80) has been observed to be impaired by these molecules, regardless of whether the individual is in a state of remission or relapse. The present theoretical framework pertaining to podocyte abnormalities and their influence on glomerular filtration in the pathogenesis of proteinuria offers an explanation for the disruption of the filtration barrier [50]. The molecules mentioned, namely sphingomyelin phosphodiesterase acid-like 3b (SMPDL3b), angiopoietin-like 4 (Angptl4), and glomerular podocyte B7-1 (CD80), have been found to interfere with the expression of these molecules. This deficiency occurs independently of an individual's status, whether in remission or relapse [51]. In accordance with previous studies, people who have nephrotic syndrome release more inflammatory cytokines, such as IL-1, IL-2, and TNF. In MCD-related nephropathy, elevated IL-13 levels are associated with proteinuria, high cholesterol levels, and podocyte fusion in feet. Immunosuppressive drugs reduce the symptoms of nephrotic syndrome, which is functionally caused by inflammatory cytokines [52].

Despite previous reports of lower SOD concentrations, this study discovered that cellular GSH and plasma levels of vitamin E decreased in patients with nephrotic syndrome. Although ceruloplasmin (CP) is a strong antioxidant and free radical scavenger, plasma CP levels increase. However, unlike earlier studies that found an increase in MDA levels in nephrotic syndrome, erythrocyte MDA, a byproduct of the lipid peroxidation

process, failed to show any appreciable alterations [53].

Research has shown that antioxidant levels are important for the prevention nephrotic syndrome. Homocysteine causes oxidative changes to LDLC and NS and affects how well steroids work [54,55].

Podocytes, a type of developing cell in the kidney basement membrane, prevent proteins from escaping in the urine by making processes which prevent protein loss. The interdiaphragm obstructs the kidneys, allowing certain substances to pass through it. FP podocytes contract and move and are held in GBM by 3/1-integrin proteins and dystroglycans. NS family gene studies have identified SD proteins, such as podocin, nephrin, -actinin-4, and TRPC6, which demonstrate how SD is placed in tandem [56].

MCD is characterised by selective albuminuria associated with FP depletion. No mutations were detected in any of these patients. This is because NS nephrins are unlikely to be produced by SD. The albumin receptor transport system moves albumin through podocytes via endocytosis and exocytosis, which explains selective albuminuria through MCS [57].

Antioxidants in nephrotic syndrome

For thousands of years, herbal medicines have been used as alternative treatments for a wide spectrum of illnesses. The antioxidative qualities of herbs and their components that come from them have been investigated for their possible therapeutic use in podocytopathies. Table 2 summarises herbs and components used for antioxidant supplementation in cases of idiopathic nephrotic syndrome, animal experimentation, and possible targeted mechanisms. These herbs and their components reduce intracellular reactive oxygen species (ROS) levels by upregulating the respiration of mitochondrial chain complex activities and downregulating pro-oxidative enzymes. In

addition, they control the ROS-mediated apoptotic pathways. Geniposide, an ingredient of *Gardenia jasminoides* Ellis, can mitigate podocyte damage and inhibit nephropathy formation [58].

Abnormalities in Cu and Zn metabolism have been documented in NS patients. Elevated homocysteine levels and low levels of B-vitamins involved in metabolism play a role in thrombosis [59]. Significant increases in serum lipid peroxidation levels of homocysteine and decreases in serum antioxidant capacity and Cu, Zn, and plasma vitamin C levels were observed in patients with NS. Antioxidant administration improves the antioxidant profile, which can play a role in preventing the appearance of kidney scar tissue [60].

In patients with renal disease, deep antioxidant therapy has the potential to decrease oxidative stress, which is a related disease. Numerous complications of nephropathy and an increase in the production of inflammatory cytokines are caused by overproduction of reactive oxygen species and free radicals in the body. Magnesium and zinc, in combination with vitamins C and E, enhance glomerular but not tubular functions [61]. Following antioxidant and mineral therapies with B-complex vitamins, significant decreases in lipid peroxide homocysteine and increases in vitamin C, total antioxidant capacity, and copper and zinc activities were noted. Research has demonstrated that B complexes, vitamins, minerals, and antioxidants can

prevent hyperhomocysteinemia and lipid peroxidation [62].

The proportion of children in the fortification group who were prehypertensive or hypertensive was significantly lower than that of the control group. Combining the antioxidants tocopherol and vitamin C can reduce the blood pressure in rats with high blood pressure. Vitamin C helps to regulate blood pressure and protects against oxidative stress by inhibiting nitric oxide synthase. In addition, vitamins C and E also promote arterial stiffness and endothelial-dependent vasodilation [46].

Another study indicated that either the breakdown of lipids or lipid peroxidation was significantly impacted by a combination of vitamins C and E. However, it was found that taking these vitamins simultaneously reduced ox-LDL and Vascular Cell Adhesion Molecule-1 (sVCAM)-1 levels in patients who had entered remission. Nitric oxide synthase has been shown to be affected by vitamin C, which reduces blood pressure, whereas vitamin E reduces blood vessel tightness and endothelial-dependent vasodilation. Overall, supplementation with vitamins C and E showed potential in reducing ox-LDL and sVCAM-1 levels during remission [63]. Our previous study showed that temulawak and black cumin extracts have lower lipid levels in NS rats. Nevertheless, there was no discernible change in HDL or LDL levels. Further research is required to fully comprehend their hypocholesterolemic qualities, more research is required [48].

TABLE 2 Herbs/components antioxidant supplementation in cases of idiopathic nephrotic syndrome

No.	Herbs/components	Animal experimentation	Targeted mechanism,	Ref.
1.	<i>Salvia przewalskii</i>	Adriamycin induced rats	Leonurine reduces proteinuria, macrophage infiltration, and early kidney damage. This could prevent oxidative stress and glomerular disorders.	[64]
2.	Berberine	Adriamycin induced rats	Berberine therapy decreases caused podocyte deaths; 4-phenylbutyric	[65]

			acid substantially decreases induced podocyte apoptosis and ER (Endoplasmic reticulum) stress-related proteins; and N-acetylcysteine reduces excessive ROS generation, perhaps defending against induced podocyte apoptosis.	
3.	<i>Ulva pertusa</i>	Adriamycin induced rats	In the kidney and blood, HU increased the activity of the antioxidant enzymes GSH-Px, CAT, and SOD. It might have done so by upregulating Nrf2 expression while reducing Keap1 expression. To sum up, HU seems to help lower oxidative stress while regulating the Keap1/Nrf2 signalling pathway, which in turn assists in reducing NS in rats.	[66]
4.	Citral	Adriamycin induced rats	The Chinese medicinal herb <i>Litsea cubeba</i> contains citral, which has the ability to reduce oxidant activity, macrophage activity, and NF- κ B activation. Citral therapy was associated with reduced inflammation, peri-glomerular inflammation, and FSGS lesions, suggesting that the drug may have renoprotective effects on the illness.	[67]
5.	<i>Epigallocatechin-3-Gallate</i>	Adriamycin induced rats	The production of ROS, TUNEL positive cells, HIF-1 α , ANGPTL4, and caspase-3 expression all appeared to be increased in animals treated with Adriamycin. Treatment with EGCG reduced these changes, with ANGPTL4, caspase-3, and Nox1 inhibition being more noticeable.	[68]
6.	Osthole	Adriamycin induced rats	Osthole stopped the acFSGS model from getting worse in mice that were given it. This was mostly because the Nrf2-mediated antioxidant system was turned on. Osthole could be a chemical that protects the kidneys because it decreased NF-B activation, COX-2 expression, PGE2 production, podocyte damage, and apoptosis.	[69]

Challenges and future development

A link exists between a decrease in organ function and a decline in kidney function, which has been linked to the breakdown of the glomerular basement membrane (GBM) and is

defined by a history of proteinuria. Edema can be induced in people by proteinuria resulting from nephrotic syndrome, isolated proteinuria, or hypertension. Owing to haemodynamic factors, filtration barrier properties, and molecular characteristics,

functional impairment in the nephrotic syndrome (NS) improves the susceptibility of the glomerular filtration barrier to materials. GFB may become increasingly permeable if disrupted, and the main problem in nephrotic syndrome (NS) is greater glomerular capillary wall permeability, which causes severe proteinuria and low albumin levels. This is because of podocyte kidney damage, which can be observed by the fact that the foot processes of podocytes in tissues from patients with NS are thinner. ROS damages cells through lipid peroxidation, which weakens the structural stability of tubular epithelial cells and makes it easier for proteins to leak out of the glomeruli. Antioxidants can improve antioxidant makeup, which can prevent scar tissue formation in the kidneys. The glomerular filtration barrier (GFB) is composed of three layers (glomerular capillary internal endothelium, glomerular visceral epithelial cells, and podocytes), and is a part of the kidney that is most affected by NS. Foot processes or peduncles of podocytes are an important part of kidney function. Ascites in the NS can be halved by eliminating ROS and inhibiting NF- κ B. This was because the changes in membrane leakage caused by protein reflection levels ceased. Further studies are warranted to identify the association between reactive oxygen species, adaptive mechanisms, and potential therapeutic strategies for reducing the burden of NS.

Conclusion

Increased sensitivity to the glomerular filtration barrier occurs with Nephrotic syndrome impairment, although the production of kidney scar tissue is prevented by antioxidants. Further investigation is required to understand the reactive oxygen species and treatment approaches.

Acknowledgements

The authors would like to thank Kusmiyati Tjahjono DK, MD, MSc, PhD, and Saphira Ayu, MD, M.Sc, for their constructive criticism of this manuscript.

Disclosure Statement

The authors declare that they have no conflicts of interest.

Authors' contributions

The authors fully accept responsibility for all the work's content and have given permission for it to be submitted.

Funding

No specific grant for this research was provided by funding organisations in the public, private, or non-profit sectors.

Data availability

Because the information in the review article was derived from the sources of the literature suggested in the article, it did not contain any associated data. This study was registered with the Open Science Framework (OSF).

Abbreviations

Angptl4: angiotensin-like 4, CHOP (C/EBP homologous protein)-TXNIP (thioredoxin-interacting protein), CMV: cytomegalovirus, CP: ceruloplasmin, C0-Q10: Coenzyme Q10, ER stress-related proteins: Endoplasmic reticulum stress related protein, FFA: Free fatty acids, FSGS: Focal segmental glomerular sclerosis, GFB: glomerular filtration barrier, GSH: glutathione, HIF-1 α : Hypoxia-inducible factor-1, IL-1: Interleukin-1, IL-2: Interleukin-2, LPL: Lipoprotein lipase, LDLc: low density lipoprotein cholesterol, SRNS: steroid-resistant nephrotic syndrome, MCD: minimal change disease, NO: nitric oxide, NPHS1: Congenital Nephrotic Syndrome of the Finnish

type, NPHS2: NPHS2 Stomatin Family Member, NS: Nephrotic syndrome, ox-LDL: oxidized LDL, PAN: Puromycin Aminonucleoside-Induced ROS: reactive oxygen species, RNS: reactive nitrogen species, SOD: super oxide dismutase, TNF: Tumor Necrotizing Factor, TUNEL: Terminal deoxynucleotidyl transferase dUTP nick end labeling, VCAM-1: Vascular Cell Adhesion Molecule-1, WT1: Wilms tumor protein gene.

Orcid:

Amallia Nuggetsiana Setyawati:

<https://orcid.org/0000-0002-1322-1369>

References

[1] (a) S.D. Meo, P. Venditti, Evolution of the knowledge of free radicals and other oxidants, *Oxid. Med. Cel. Longev.*, **2020**, 2020. [Crossref], [Google Scholar], [Publisher], (b) R. Miftari, H. Maloku, V. Topciu, A. Bajqinca, Y. Kaciu, M. Shabani, M. Ukimeraj, B. Maloku, Congenital anomalies of urinary tract in Kosovo population diagnosed by renal scintigraphy, *J. Med. Chem. Sci.*, **2023**; 6, 1457-1468. [Crossref], [Google Scholar], [Publisher], (c) M. Hashemzadeh Esfahani, B. Haghighi Aski, A. Manafi Anari, G. Gharehbaghi, M. Sakhaei, E. Shorafa, N. Hooman, Extreme hypertension associated with SARS-COVID-19 infection in pediatrics with kidney disease: a retrospective observational study, *J. Med. Chem. Sci.*, **2023**; 6, 79-85. [Crossref], [Google Scholar], [Publisher]

[2] H. Sies, D.P. Jones, Reactive oxygen species (ROS) as pleiotropic physiological signalling agents, *Nat. Rev. Mol. Cell Biol.*, **2020**, 21, 363-383. [Crossref], [Google Scholar], [Publisher]

[3] (a) M. Sharifi-Rad, N.V. Anil Kumar, P. Zucca, E. Maria Varoni, L. Dini, E. Panzarini, J. Rajkovic, P.V. Tsouh Fokou, E. Azzini, I. Peluso, A. Prakash Mishra, M. Nigam, Y. El Rayess, M. El Beyrouthy, L. Polito, M. Iriti, N. Martins, M. Martorell, A.O. Docea, W.N. Setzer, D. Calina, W.C. Cho, J. Sharifi-Rad, Lifestyle, Oxidative stress, and antioxidants: back and forth in the

pathophysiology of chronic diseases, *Front. Physiol.*, **2020**, 2, 694. [Crossref], [Google Scholar], [Publisher], (b) B. Korir, J. Kibet, B. Mosonik, A review of the current trends in the production and consumption of bioenergy, *Prog. Chem. Biochem. Res.*, **2021**, 4, 109-133. [Crossref], [Google Scholar], [Publisher], (c) O. Solomon, W. Rabiou Saidou Umar, H. Sanusi Wara, A. Sadiq Yakubu, M. Michael Azubuike, M. Asugu Mary, H. Louis, Antiulcerogenic activity of methanol extract and solvent fractions of Stem Bark of *Lannea acida* (A. Rich) against ethanol-induced gastric mucosal injury in albino rats, *Prog. Chem. Biochem. Res.*, **2018**, 1, 29-39. [Crossref], [Google Scholar], [Publisher], (d) G. Chala, Review on green synthesis of iron-based nanoparticles for environmental applications, *J. Chem. Rev.*, **2023**, 5, 1-14. [Crossref], [Google Scholar], [Publisher], (e) N. Vaeli, Laboratory study of effective factors on how to extract carvacrol from *oliveria decumbens* plant with the help of supercritical fluid CO₂ and using ultrasound waves, *Eurasian J. Sci. Technol.*, **2022**; 2, 32-43. [Crossref], [Google Scholar], [Publisher], (f) Z.A. Messaoudi, D. Lahcene, T. Benaissa, M. Messaoudi, B. Zahraoui, M. Belhachemi, A. Choukchou-Braham, Adsorption and photocatalytic degradation of crystal violet dye under sunlight irradiation using natural and modified clays by zinc oxide, *Chem. Methodol.*, **2022**; 6, 661-676. [Crossref], [Google Scholar], [Publisher], (g) B. Bulus, Applications of *justicia secunda* extracts in functional foods and natural products: a review, *Adv. J. Chem. Sect. B. Nat. Prod. Med. Chem.*, **2023**; In Press, 1-10. [Crossref], [Pdf], [Publisher]

[4] R. Sotler, B. Poljšak, R. Dahmane, T. Jukić, D. Pavan Jukić, C. Rotim, P. Trebše, A. Starc, Prooxidant activities of antioxidants and their impact on health, *Acta Clin Croat.*, **2019**, 58, 726-736. [Crossref], [Google Scholar], [Publisher]

[5] X. Zhang, J. Xu, H. Xiao, Y. Yao, H. Wang, Y. Ren, M. Liu, F. Wang, X. Zhong, X. Liu, B. Su, M. Cheng, L. Chai, J. Ding, S. Wang, Value of electron microscopy in the pathological diagnosis of native kidney biopsies in children,

- Pediatr Nephrol.*, **2020**, *35*, 2285-2295. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [6] L.P.E.S Kresnandari, G.A.P Nilawati, I.G.A.T Windiani, I.G.A.N.S Adnyana, Soetjningsih, B.N Mahakrishna, N.L.S.P. Murti, The quality of life in children with nephrotic syndrome at Prof IGNG Ngoerah Hospital, Denpasar, *Bali. Intisari Sains Medis.*, **2023**, *14*, 401-406. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [7] A. Trautmann, M. Vivarelli, S. Samuel, D. Gipson, A. Sinha, F. Schaefer, N. Kar Hui, O. Boyer, M.A. Saleem, L. Feltran, J. Müller-Deile, J.U. Becker, F. Cano, H. Xu, Y. Ngo Lim, W. Smoyer, I. Anochie, K. Nakanishi, E. Hodson, D. Haffner, IPNA clinical practice recommendations for the diagnosis and management of children with steroid-resistant nephrotic syndrome, *Pediatr Nephrol.*, **2020**, *35*, 1529-1561. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [8] A.N. Setyawati, O. Mellyana, H. Muryawan, Medication adherence in patients with nephrotic syndrome., *Pak. J. Med. Health Sci.*, **2020**, *14*, 411-414. [[Publisher](#)]
- [9] Y.L. Lv, N. Guan, J. Ding, Y. Yao, H.J. Xiao, X.H. Zhong, F. Wang, X.Y. Liu, H.W. Zhang, B.G. Su, K. Xu, Spectrum of thrombotic complications and their outcomes in Chinese children with primary nephrotic syndrome, *Ital. J. Pediatr.*, **2020**, *46*, 1-8. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [10] S Lionaki, G. Liapis, J.N. Boletis, Pathogenesis and management of acute kidney injury in patients with Nephrotic syndrome due to primary glomerulopathies, *Medicina*, **2019**, *55*. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [11] V. Tesar, T. Zima, M. Kalousová, Pathobiochemistry of nephrotic syndrome, *Adv Clin Chem.*, **2003**, *37*, 173-218. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [12] J. Tecklenborg, D. Clayton, S. Siebert, S.M. Coley, The role of the immune system in kidney disease, *Clin Exp Immunol.*, **2018**, *192*, 142-150. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [13] A. Abid, S. Shahid, M. Shakoor, A.A. Lanewala, S. Hashmi, S. Khaliq, Screening of the LAMB2, WT1, NPHS1, and NPHS2 genes in pediatric nephrotic syndrome, *Front Genet.*, **2018**, *9*, 214. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [14] H.H. Haeri, J. Eisermann, H. Schimm, A. Büscher, P. Hoyer, D. Hinderberger, Profound changes in functional structure and dynamics of serum albumin in children with nephrotic syndrome: an exploratory research study, *J Med Chem.*, **2023**, *66*, 12115-12129. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [15] T. Nishimura, O. Uemura, S. Hibino, K. Tanaka, R. Kitagata, S. Yuzawa, T. Kasagi, N. Fujita, Serum albumin level is associated with mycophenolic acid concentration in children with idiopathic nephrotic syndrome, *Eur J Pediatr.*, **2022**, *181*, 1159-1165 [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [16] E.M. Yang, K.H. Yoo, Y.H. Ahn, S.H. Kim, J.W. Lee, W.Y. Chung, M.H. Cho, K.H. Kim, H.Cho, M.J. Lee, J.S. Suh, H.S. Hyun, J.M. Lee, M. H. Cho, J.H. Kim, I.S. Ha, H. Cheong, H.G. Kang, Lower albumin level and longer disease duration are risk factors of acute kidney injury in hospitalized children with nephrotic syndrome, *Pediatr. Nephrol.*, **2021**, *36*, 701-709. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [17] P.B. Soeters, R.R. Wolfe, A. Shenkin, Hypoalbuminemia: Pathogenesis and Clinical Significance, *JPEN J Parenter Enteral Nutr.*, **2019**, *43*, 181-193 [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [18] A.A. Jackson, Albumin in nephrotic syndrome and oedematous malnutrition, *Paediatr. Int. Child Health*, **2015**, *35*, 77-80. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [19] J. Meena, A. Bagga, Current Perspectives in Management of Edema in Nephrotic Syndrome, *Indian J Pediatr.*, **2020**, *87*, 633-640. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [20] P. Claudio, M. Gabriella, Nephrotic syndrome: pathophysiology and consequences, *J Nephrol.*, **2023**, *19*. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]

- [21] B.A. Molitoris, R.M. Sandoval, S.P.S. Yadav, M.C. Wagner, Albumin uptake and processing by the proximal tubule: physiological, pathological, and therapeutic implications, *Physiol Rev.*, **2022**, *102*, 1625-1667. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [22] A. Edwards, K.R. Long, C.J. Baty, K.E. Shipman, O.A. Weisz, Modelling normal and nephrotic axial uptake of albumin and other filtered proteins along the proximal tubule, *J Physiol.*, **2022**, *600*, 1933-1952. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [23] W.D. Comper, J. Vuchkova, K.J. McCarthy, New insights into proteinuria/albuminuria, *Frontiers in Physiology*, **2022**, *13*, 991756. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [24] S. Hatakeyama, A. Tojo, H. Satonaka, N.O. Yamada, T. Senda, T. Ishimitsu, Decreased Podocyte Vesicle Transcytosis and Albuminuria in APC C-Terminal Deficiency Mice with Puromycin-Induced Nephrotic Syndrome, *Int. J. Mol. Sci.*, **2021**, *22*, 13412. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [25] A. Matyjek, S. Literacki, S. Niemczyk, A. Rymarz, Protein energy-wasting associated with nephrotic syndrome - the comparison of metabolic pattern in severe nephrosis to different stages of chronic kidney disease, *BMC Nephrol.*, **2020**, *21*, 346. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [26] J.I. Maier, M. Rogg, M. Helmstädter, A. Sammarco, G. Walz, M. Werner, C. Schell, A Novel Model for Nephrotic Syndrome Reveals Associated Dysbiosis of the Gut Microbiome and Extramedullary Hematopoiesis, *Cells.*, **2021**, *10*, 1509. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [27] E.C. Ray, H. Rondon-Berrios, C.R. Boyd, T.R. Kleyman, Sodium retention and volume expansion in nephrotic syndrome: implications for hypertension, *Adv Chronic Kidney Dis.*, **2015**, *22*, 179-84. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [28] S. Agrawal, J.J. Zaritsky, A. Fornoni, W.E. Smoyer, Dyslipidaemia in nephrotic syndrome: mechanisms and treatment, *Nat Rev Nephrol.*, **2018**, *14*, 57-70. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [29] K.J. Hampson, M.L. Gay, M.E. Band, Pediatric Nephrotic Syndrome: Pharmacologic and Nutrition Management, *Nutr Clin Pract.*, **2021**, *36*, 331-343. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [30] S. Verma, P. Singh, S. Khurana, N.K. Ganguly, R. Kukreti, L. Saso, D.S. Rana, V. Taneja, V. Bhargava, Implications of oxidative stress in chronic kidney disease: a review on current concepts and therapies, *Kidney Res Clin Pract.*, **2021**, *40*, 183-193. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [31] D.M. Small, J.S. Coombes, N. Bennett, D.W. Johnson, G.C. Gobe, Oxidative stress, antioxidant therapies and chronic kidney disease, *Nephrology (Carlton)*, **2012**, *17*, 311-21. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [32] B.M. Lane, M. Chryst-Stangl, G. Wu, M. Shalaby, S. El Desoky, C.C. Middleton, K. Huggins, A. Sood, A. Ochoa, A.F. Malone, R. Vancini, S.E. Miller, G. Hall, S.Y. Kim, D.N. Howell, J.A. Kari, R. Gbadegesin, Corresponding, Steroid-sensitive nephrotic syndrome candidate gene CLVS1 regulates podocyte oxidative stress and endocytosis, *JCI Insight.*, **2022**, *7*, e152102. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [33] M.K. Foret, R. Lincoln, S. Do Carmo, A.C. Cuello, G. Cosa, Connecting the "Dots": from free radical lipid autoxidation to cell pathology and disease, *Chem Rev.*, **2020**, *120*, 12757-12787. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [34] K.N. Moorani, K.M. Khan, A. Ramzan, Infections in children with nephrotic syndrome, *J Coll Physicians Surg Pak.*, **2003**, *13*, 337-9. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [35] H.J. Forman, H. Zhang, Targeting oxidative stress in disease: promise and limitations of antioxidant therapy, *Nat Rev Drug Discov.*, **2021**, *20*, 689-709. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [36] Q. Ye, D.J. Wang, B. Lan, J.H. Mao, T-cell and B-cell repertoire diversity are selectively

- skewed in children with idiopathic nephrotic syndrome revealed by high-throughput sequencing, *World J. Pediatr.*, **2023**, *19*, 273-282. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [37] J.B. Kopp, H.J. Anders, K. Susztak, M.A. Podestà, G. Remuzzi, F. Hildebrandt, P. Romagnani. Podocytopathies, *Nat Rev Dis Primers.*, **2020**, *6*, 68. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [38] T. Ozeki, S. Maruyama, T. Imasawa, T. Kawaguchi, H. Kitamura, M. Kadomura, R. Katafuchi, K. Oka, H. Yokoyama, H. Sugiyama, H. Sato, Clinical manifestations of focal segmental glomerulosclerosis in Japan from the Japan Renal Biopsy Registry: age stratification and comparison with minimal change disease, *Sci Rep.*, **2021**, *11*, 2602. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [39] N. Pallet, A.A. Fernández-Ramos, M.A. Lorient, Impact of Immunosuppressive Drugs on the Metabolism of T Cells, *Int Rev Cell Mol Biol.*, **2018**, *341*, 169-200. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [40] S.J. Park, Y. Kim, C. Li, J. Suh, J. Sivapackiam, T.M. Goncalves, G. Jarad, G. Zhao, F. Urano, V. Sharma, Y.M. Chen, Blocking CHOP-dependent TXNIP shuttling to mitochondria attenuates albuminuria and mitigates kidney injury in nephrotic syndrome, *Proc Natl Acad Sci USA.*, **2022**, *119*, e2116505119. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [41] W. Tan, R. Airik, Primary coenzyme Q10 nephropathy, a potentially treatable form of steroid-resistant nephrotic syndrome, *Pediatr Nephrol.*, **2021**, *36*, 3515-3527. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [42] G. Kleiner, E. Barca, M. Ziosi, V. Emmanuele, Y. Xu, A. Hidalgo-Gutierrez, C. Qiao, S. Tadesse, E. Area-Gomez, L.C. Lopez, C.M. Quinzii, CoQ10 supplementation rescues nephrotic syndrome through normalization of H2S oxidation pathway, *Biochim Biophys Acta Mol. Basis Dis.*, **2018**, *1864*, 3708-3722. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [43] S. Drovandi, B.S. Lipska-Ziętkiewicz, F. Ozaltin, F. Emma, B. Gulhan, O. Boyer, A. Trautmann, H. Xu, Q. Shen, J. Rao, K.M. Riedhammer, U. Heemann, J. Hoefele, S.L. Stenton, A.N. Tsygin, K.H. Ng, S. Fomina, E. Benetti, M. Aurelle, L. Prikhodina, M.F. Schreuder, M. Tabatabaieifar, M. Jankowski, S. Baiko, J. Mao, C. Feng, C. Liu, S. Sun, F. Deng, X. Wang, S. Clavé, M. Stańczyk, I. Bałasz-Chmielewska, M. Fila, A.M. Durkan, T.K. Levart, I. Dursun, N. Esfandiar, D. Haas, A. Bjerre, A. Anarat, M.R. Benz, S. Talebi, N. Hooman, G. Ariceta, PodoNet Consortium; mitoNET Consortium; CCGKDD Consortium; Schaefer F. Oral Coenzyme Q10 supplementation leads to better preservation of kidney function in steroid-resistant nephrotic syndrome due to primary Coenzyme Q10 deficiency, *Kidney Int.*, **2022**, *102*, 604-612. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [44] H. Sadeghi, M. Mansourian, E. Panahi kokhdan, Z. Salehpour, I. Sadati, Kazem A-Goudarzi, A. Asfaram, A.H. Doustimotlagh, Antioxidant and protective effect of *Stachys pilifera* Benth against nephrotoxicity induced by cisplatin in rats, *J Food Biochem.*, **2020**, *5*, e13190. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [45] F. Hashmi, S.S. Ghazi, H. Mumtaz, M Amjad, N Zaman, S Ahmad, Role of zinc supplementation in reducing relapses in steroid sensitive nephrotic syndrome, *Annals of PIMS-Shaheed Zulfiqar Ali Bhutto Medical University*, **2021**, *17*, 236-240. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [46] O. Mellyana, R. Widajat, N. Sekarwana, Combined supplementation with α -tocopherol and vitamin C improves the blood pressure of pediatric idiopathic nephrotic syndrome patients, *Clin. Nut. Exp.*, **2019**, *23*, 16-121. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [47] H.Y. Fan, X.K. Wang, X. Li, K. Ji, S.H. Du, Y. Liu, L.L. Kong, J-c. Xu, G-q. Yang, D-q. Chen, D. Qi, Curcumin, as a pleiotropic agent, improves doxorubicin-induced nephrotic syndrome in rats, *J Ethnopharmacol.*, **2020**, *250*, 112502. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]

- [48] J.R. Hadi, K. Tjahjono, R. Halleyantoro, P.K. Dewi, F. Fulyani, A.N. Setyawati, Combination of *Nigella sativa* and *Curcuma xanthorrhiza* Roxb, reduces hyperlipidemia in Nephrotic syndrome, *J Pharm Negat Results*, **2023**, 1804-1812. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [49] T. Liu, L. Zhang, D. Joo, S.C. Sun, NF- κ B signaling in inflammation, *Signal Transduct Target Ther.*, **2017**, *2*, 17023. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [50] S. Watanabe, K. Hirono, T. Aizawa, K. Tsugawa, K. Joh, T. Imaizumi, H. Tanaka, Podocyte sphingomyelin phosphodiesterase acid-like 3b decreases among children with idiopathic nephrotic syndrome, *Clin Exp Nephrol.*, **2020**, *25*. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [51] S. Watanabe, U. Hidenori, S Hashimoto, S. Riko, T. Aizawa, K. Tsugawa, T. Imaizumi, H. Tanaka. Sphingomyelin phosphodiesterase acid-like 3b is essential for toll-like receptor 3 signaling in human podocytes, *The J. Membr. Biol.*, **2022**, *255*, 117-122. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [52] E. Ahmadian, Y.R. Saadat, E.D. Abdolahinia, M. Bastami, M.M. Shoja, S. Zununi Vahed, M. Ardalani, The Role of cytokines in nephrotic syndrome, *Mediators Inflamm*, **2022**, *2022*, 6499668. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [53] M. Khurana, D.M. Silverstein, Etiology and management of dyslipidemia in children with chronic kidney disease and end-stage renal disease, *Pediatr Nephrol.*, **2015**, *30*, 2073-84. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [54] O.P. Mishra, A.K. Gupta, R. Prasad, Z. Ali, R.S. Upadhyay, S.P. Mishra, N.K. Tiwary, F.S. Schaefer, Antioxidant status of children with idiopathic nephrotic syndrome, *Pediatr Nephrol.*, **2011**, *26*, 251-6. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [55] K. Udwan, G. Brideau, M. Fila, A. Edwards, B. Vogt, A. Doucet, Oxidative stress and nuclear factor κ B (NF- κ B) increase peritoneal filtration and contribute to ascites formation in nephrotic syndrome, *J. Biol. Chem.*, **2016**, *291*, 11105-13. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [56] B.E. Drummond, W.S. Ercanbrack, Drand R.A. Wingert, Modeling Podocyte Ontogeny and Podocytopathies with the Zebrafish, *J Dev Biol*, **2023**, *11*. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [57] L. Butt, D. Unnersjö-Jess, M. Höhne, A. Edwards, J. Binz-Lotter, D. Reilly, R. Hahnfeldt, V. Ziegler, K. Fremter, M.M. Rinschen, M. Helmstädter, L.K. Ebert, H. Castrop, M.J. Hackl, G. Walz, P.T. Brinkkoetter, M.C. Liebau, K. Tory, P.F. Hoyer, B.B. Beck, H. Brismar, H. Blom, B. Schermer, T. Benzing, A molecular mechanism explaining albuminuria in kidney disease, *Nat. Metab.*, **2020**, *2*, 461-474. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [58] Y.T. Zhu, C. Wan, J.H. Lin, H.P. Hammes, C. Zhang, Mitochondrial oxidative stress and cell death in podocytopathies, *Biomolecules*, **2022**, *12*, 403. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [59] K. Jomova, M. Valko, Advances in metal-induced oxidative stress and human disease. *Toxicology*, **2011**, *283*, 65-87. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [60] J. Dwivedi, P.D. Sarkar, Interrelationship of oxidative stress, homocysteine, lipoprotein (a), copper and zinc in Nephrotic syndrome: recent advances: risk for cardiac diseases, *Pharmacologyonline*, **2009**, *2*, 605-5. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [61] S.F. Rapa, B.R. Di Iorio, P. Campiglia, A. Heidland, S. Marzocco, Inflammation and oxidative stress in chronic kidney disease-potential therapeutic role of minerals, vitamins and plant-derived metabolites, *Int J Mol Sci.*, **2019**, *21*, 263. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [62] W. Hamik, D. Hilmanto, S. Rahayuningsih, Relationship between serum zinc and homocysteine in children with nephrotic syndrome, *Pl.*, **2019**, *59*, 98-03. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]

- [63] O. Mellyana, E. Dharmana, H. Susanto, N. Sekarwana, Effects of combined supplementation of vitamins C and E on the oxidative modification of low-density lipoprotein, soluble form of CD36, soluble vascular cell adhesion molecule-1, and nitrite/nitrate oxide levels in idiopathic nephrotic syndrome, *Biomark. Genom. Med.*, **2015**, *7*, 125-130. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [64] X. Liu, W. Cao, J. Qi, Q. Li, M. Zhao, Z. Chen, J. Zhu, Z. Huang, L. Wu, B. Zhang, Y. Yuan, C. Xing, Leonurine ameliorates adriamycin-induced podocyte injury via suppression of oxidative stress, *Free Radic Res.*, **2018**, *52*, 952-960. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [65] X.Y. Xiang, T. Liu, Y. Wu, X.-S. Jiang, J.-L. He, X.-M. Chen, X.-G. Du, Berberine alleviates palmitic acid-induced podocyte apoptosis by reducing reactive oxygen species-mediated endoplasmic reticulum stress, *Mol Med Rep.*, **2021**, *23*, 3. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [66] Y. Wan, S. Wang, K. Chen, L. Liu, X. Wang, B. Zhang, L. Hu, S. Liu, T. Zhao, H. Qi, High-sulfated derivative of polysaccharide from *Ulva pertusa* improves Adriamycin-induced nephrotic syndrome by suppressing oxidative stress, *Food Funct.*, **2023**. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [67] S.M. Yang, K.F. Hua, Y.C. Lin, A. Chen, J.M. Chang, L.K. Chao, C.L. Ho, S.M. Ka. Citral is renoprotective for focal segmental glomerulosclerosis by inhibiting oxidative stress and apoptosis and activating Nrf2 pathway in mice, *PLoS One.*, **2013**, *8*, e74871. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [68] G. Liu, L. He, Epigallocatechin-3-Gallate Attenuates Adriamycin-Induced Focal Segmental Glomerulosclerosis via Suppression of Oxidant Stress and Apoptosis by Targeting Hypoxia-Inducible Factor-1 α /Angiopoietin-Like 4 Pathway, *Pharmacology*, **2019**, *103*, 303-314. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [69] S.M. Yang, Y.L. Chan, K.F. Hua, J.M. Chang, H.L. Chen, Y.J. Tsai, S.H. Wu, Y.-F. Wang, C.-L. Tsai, A. Chen, S.M. Ka, Osthole improves an accelerated focal segmental glomerulosclerosis model in the early stage by activating the Nrf2 antioxidant pathway and subsequently inhibiting NF- κ B-mediated COX-2 expression and apoptosis, *Free Radic. Biol. Med.*, **2014**, *73*, 260-269. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]

How to cite this article: Amallia Nuggettsiana Setyawati, The role of oxidative stress in hypoalbuminemia nephropathy related to Nephrotic syndrome: a critical review. *Eurasian Chemical Communications*, 2024, 6(1), 32-49. **Link:** https://www.echemcom.com/article_182691.html